

## **Influence of Crude Oil Contamination on the Chemical and Microbiological Aspects of Saudi Arabian Soils**

**A.R. Hashem**

*Department of Botany and Microbiology, College of Science,  
King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia*

(Received on 3 January 1994; accepted for publication on 15 January 1995)

**Abstract.** Samples of petroleum-contaminated and uncontaminated soils were collected from Eastern region of Saudi Arabia and analyzed for soil texture, organic matter content, total soluble salts, pH, mineral content, fungal and bacterial microflora as indicators of soil pollution. The results obtained revealed that both petroleum-contaminated and uncontaminated soils were sandy, mildly alkaline, containing almost the same amounts of organic matter, soluble salts and heavy minerals. The total number of colonies of bacteria and fungi per gram of soil were higher in uncontaminated than petroleum contaminated soils. A total number of twenty fungal species were isolated from both contaminated and uncontaminated soils. The uncontaminated soil is capable of supporting the growth of all 20 species while the contaminated soil supports only 17 of these species. In both contaminated and uncontaminated soils *Aspergillus* was the dominant fungal genus (6 species) followed by *Penicillium* and *Cladosporium* (3 species each). Bacterial genera belonging to *Arthrobaacter*, *Bacillus*, *Micrococcus*, *Pseudomonas* and *Staphylococcus* were also isolated.

### **Introduction**

Most of the Eastern region of Saudi Arabia including coastal area, bordering Kuwait, became polluted with petroleum during the recent Gulf war. Microorganisms including bacteria and fungi could degrade petroleum as a carbon and energy source [1, 2] [3, p. 692], [4-7].

*Aspergillus*, *Penicillium*, *Trichoderma*, and *Cladosporium* species could degrade petroleum product and other high carbon compounds [8, 9, 10]. Species of bacterial genera like, *Pseudomonas*, *Bacillus*, *Arthrobaacter*, *Micrococcus*, and *Staphylococcus* could also degrade petroleum products and other high carbon organic compounds [6, 10, 11]. Edible oils, petroleum products, emulsion paints, diesel fuels etc. could be converted into useful products by biodegradation activity of these microorganisms [12, 13].

There are several problems dealing with petroleum contaminated soils which effect human health and environmental quality [14]. This study considers the influence of

spilled oil on soil chemical properties and proliferation of soil microflora in the sandy sites of Eastern Saudi Arabia.

### Materials and Methods

During the Gulf war, due to spilling of petroleum oil on the soil of the war zone bordering Kuwait, several patches of petroleum oil contamination could be seen. Samples were collected from these oil patches and also from nearby non-contaminated soil. Five samples (five replicates of each) were collected from five localities of the same area (Al-Khafji area) both from contaminated and uncontaminated soil. Total number of bacteria and fungi per gram of soil were counted by using the dilution plate method for each type of sample. These samples were then mixed together and random five gram subsamples of soil were taken and homogenized into 45 ml of presterilized water. The resulting suspension was used to isolate fungi and bacteria by the dilution plate method as described earlier [15].

For fungal isolation, Czapek's Dox agar (Oxoid Ltd., London) with rose bengal (0.033 g/L) and streptomycin sulphate (0.033 g/L) was used. For bacterial isolation, nutrient agar (Oxoid Ltd., London) was used. Identification of fungi was carried out according to Ellis [6, pp. 309, 315, 465, 498, 547], Ellis [17, pp. 414, 430], Gilman [18, pp. 170, 212, 220, 225, 334, 374], Ramirez [19, pp. 300, 471], and Raper and Fennell [20, pp. 203, 301, 357, 377, 412]. The identification of bacterial flora was done according to Buchanan and Gibbons [21].

Total soluble salts, organic matter, and soil reaction (pH value) were determined according to Bower and Wilcox [22, pp. 933-951], Broadbent [23, pp. 1397-1400], and Peech [24, pp. 920-923]. Soil texture was determined by the hydrometer method [25, pp. 77-80]. Total metal content was determined according to Hashem, [26].

Table 1. Physical and chemical analysis of soil samples

Soil treatment	Soil texture	Organic matter %	Total soluble salts %	pH value	Total metal content ( $\mu\text{g/g}$ )					
					AL	Cd	Cu	Fe	Pb	Zn
Contaminated	Sandy	1.3	1.1	7.0	90 $\pm$ 3	6 $\pm$ 0.1	13 $\pm$ 1	26 $\pm$ 2	4 $\pm$ 1	16 $\pm$ 1
Uncontaminated	Sandy	1.8	1.0	7.5	85 $\pm$ 03	4 $\pm$ 03	11 $\pm$ 2	24 $\pm$ 2	3 $\pm$ 0.1	14 $\pm$ 1

n = 5

$\pm$  standard deviation

## Results and Discussion

The results of physical and chemical analysis of soils are shown in Table 1. Both contaminated and uncontaminated soils were similar in their texture, organic matter content, total soluble salt content, pH value and total metal content. This is consistent with the previous findings by the author in Saudi Arabian soils [27, 28]. The total number of fungal and bacterial colonies per gram of uncontaminated soil is shown in Table 2. The average number of fungal and bacterial colonies per gram soil isolated from uncontaminated and contaminated soil were 2610 fungal and  $4.1 \times 10^6$  bacterial and 1280 fungal and  $4.1 \times 10^4$  bacterial respectively. The contaminated soil supports a lower number of microflora than the uncontaminated one (Tables 2 and 3).

Table 2. Number of fungal and bacterial colonies per gram of uncontaminated soil

Soil sample delineation	No of colonies per gram of soil	
	Fungi	Bacteria
A	$\pm 1.6$	$\times 10^6$
B	$\pm 14$	$\times 10^6$
C	$\pm 15$	$\times 10^6$
D	$\pm 10$	$\times 10^6$
E	$\pm 11$	$\times 10^6$
Average		$\times 10^6$

Table 3. Number of fungal and bacterial colonies per gram of contaminated soil

Soil sample delineation	No of colonies per gram of soil	
	Fungi	Bacteria
A - 1	$1260 \pm 12$	$3 \times 10^4$
B - 1	$1631 \pm 10$	$6 \times 10^4$
C - 1	$993 \pm 9$	$5.5 \times 10^4$
D - 1	$1350 \pm 14$	$3.4 \times 10^4$
E - 1	$1169 \pm 13$	$2.6 \times 10^4$
Average	1280	$4.1 \times 10^4$

The uncontaminated soil supports the growth of all of the 20 fungal species while the contaminated soil supports the growth of only 17 species out of the 20 species identified (Table 4). The dominant genera were *Aspergillus* (6 species) followed by *Cladosporium* and *Penicillium* (3 species each), *Ulocladium* and *Alternata* (2 species each).

**Table 4. Fungal flora isolated from petroleum- contaminated and uncontaminated soil**

Fungi	No of colonies per gram of soil	
	Uncontaminated	Contaminated
<i>Alternaria alternata</i>	36 ± 5	24 ± 4
<i>A. chlamydospora</i>	49 ± 6	39 ± 5
<i>Aspergillus carbonarius</i>	34 ± 7	18 ± 3
<i>A. flavus</i>	52 ± 9	29 ± 5
<i>A. funiculosus</i>	21 ± 3	14 ± 3
<i>A. niger</i>	41 ± 5	21 ± 4
<i>A. thomii</i>	16 ± 3	NF
<i>A. zonatus</i>	12 ± 3	21 ± 4
<i>Chaetomium piluliferum</i>	21 ± 4	26 ± 4
<i>Cladosporium herbarum</i>	34 ± 3	24 ± 3
<i>C. resinae</i>	26 ± 4	59 ± 6
<i>C. sphaerospermum</i>	26 ± 3	16 ± 3
<i>Eurotium</i> sp.	21 ± 5	56 ± 7
<i>Fusarium solani</i>	34 ± 4	18 ± 3
<i>Penicillium chrysogenum</i>	56 ± 3	42 ± 6
<i>P. expansum</i>	18 ± 4	NF
<i>P. notatum</i>	23 ± 4	NF
<i>Trichoderma</i> sp.	26 ± 3	46 ± 5
<i>Ulocladium atrum</i>	39 ± 5	46 ± 5
<i>U. chlamydosporum</i>	32 ± 3	39 ± 4
No. of species/genera	20/9	17/9

NF = Not found

*Aspergillus thomii*, *Penicillium expansum* and *P. notatum* were not isolated from contaminated soil. Generally, the number of colonies of a particular species per gram of soil was less in contaminated than in uncontaminated soil except for *Aspergillus zonatus*, *Cladosporium resinae*, *Eurotium* sp., *Trichoderma* sp., *Ulocladium atrum* and *U. chlamydosporum* where a reverse trend was observed.

The bacterial species *Arthrobacter*, *Bacillus*, *Micrococcus*, *Pseudomonas* and *Staphylococcus* were isolated from both uncontaminated and contaminated soil (Table 5) but the number of colonies per gram of soil were less in petroleum contaminated than in uncontaminated soil.

**Table 5. Bacteria flora isolated from petroleum- contaminated and uncontaminated soil**

Bacteria	No of colonies per gram of soil	
	Uncontaminated	Contaminated
<i>Arthrobacter</i> sp.	3539 ± 25	1696 ± 16
<i>Bacillus</i> sp.	6344 ± 26	3439 ± 15
<i>Micrococcus</i> sp.	2691 ± 21	1329 ± 14
<i>Pseudomonas</i> sp.	5648 ± 18	3126 ± 16
<i>Staphylococcus</i> sp.	3636 ± 16	1346 ± 21

*Aspergillus flavus*, *A. niger*, *Penicillium chrysogenum*, *Trichoderma* and *Ulocladium atrum* isolated here, were previously reported in petroleum contaminated soils in Saudi Arabia [29]. However, *Aspergillus ellipticus*, *Chaetomium bostrychodes*, *Mucor* and *Rhizopus* isolated earlier [29] were not encountered here. All fungal genera and species isolated in this work have previously been reported as biodegraders of various complex organic compounds and petroleum products [4, 6, 8, 9, 30, 31, 32].

The contamination of soil with toxic environmental contaminants such as petroleum is pervasive problem of potential human health concern particularly to those working and residing near hazardous products (e.g., polyaromatic hydrocarbons, benzene, lead, ...etc.) are compounds commonly identified as soil contaminants at various hazardous waste sites, and the toxicological profiles of several of these substances have already been drafted [33]. It is worth mentioning that more research is warranted to elucidate the oil contamination problem of soils encountered in spillage areas.

**Acknowledgement.** I would like to thank Mr. D. Baidridge, San Diego Sate University, U.S.A., for his valuable suggestions and assistance.

### References

- [1] Cobet, A.B.; Guard, H.E., and Chatigny, M.A. 'Considerations in Applications of Microorganisms to the Environment for Degradation of Petroleum Products.' In: *The Microbiological Degradation of Oil Pollutants*. Ahearn, D.G. and Meyers, S.P. (Eds.), pp. 81-87. Baton Rouge: Louisiana State University Press, 1973.
- [2] Davies, J.S. and Westlake, D.W.S. "Crude Oil Utilization by Fungi." *Canadian Journal of Microbiology* 25 (1979), 146-156.
- [3] Atlas, R.M. *Petroleum Microbiology*. New York: MacMillan, 1984.

- [4] Bossert, I. and Bartha, R. "The Fate of Petroleum in Soil Ecosystem." In: *Petroleum Microbiology* (R.M. Atlas, ed), New York: MacMillan, 1984.
- [5] Kirk, P.W. and Gordon, A.S. "Hydrocarbon Degradation by Filamentary Marine Higher Fungi." *Mycologia*, 80, No. 6 (1988), 776-782.
- [6] Ehrhardt, H.M. and Rehm, H. "Semicontinuous and Continuous Degradation of Phenol by *Pseudomas putida* Adsorbed on Activated Carbon." *Applied Microbiol Biotechnology* 30 (1989), 312-317.
- [7] Kref, J.U. and Schink, B. "Demethylation and Degradation of Phenylmethylethers by the Sulfide-methylation Homoacetogenic Bacterium Strain TMBS4." *Arcg. Microbiol.* 159 (1993), 308-315.
- [8] Llanos, C. and Kjoller, A. "Changes in the Flora of Soil Fungi: Following Oil Waste Application." *Oikos*, 27 (1976), 331-382.
- [9] Savitha, J. "Utilization of Hydrocarbon in Crude Oil by Fungi." *Current Science*, 2 (1986), 1248-1249.
- [10] Knoll, G. and Winter, J. "Anaerobic Degradation of Phenol in Sewage Sludge. Benzuate Formation from Phenol and Carbon Dioxide in the Presence of Hydrogen." *Applied Microbiol Biotechnology*, 25 (1989), 384-391.
- [11] Bettman, H. and Rehm, H. "Degradation of Phenol by Polymer and Entrapped Microorganisms." *Applied Microbiol Biotechnology*, 20 (1984), 285-290.
- [12] Carrod, P.A. "Enzymes and Microorganisms in Food Industry Waste Processing and Conversion to Useful Products: A Review of the Literature." *Research on Reconversion and Conservation*, 3 (1978), 165-178.
- [13] Tan, T. and Leong, W. "Screening for Extracellular Enzymes of Fungi from Manufacturing Wastes." *MIRCEN Journal*, 2 (1986), 445-452.
- [14] Tucker, R.A. "Problems Dealing with Petroleum Contaminated Soils." In: *Petroleum Contaminated Soils: Kostecki, P.T. and Calabrese, E.J. (Eds.) Vol. 1, pp. 37-54. Michigan: Lewis Publishers, Inc., 1990.*
- [15] Hashem, A.R. "Studies on the Fungal Flora of Saudi Arabian Soils." *Cryptogamic Botany*, 2, No. 3 (1991), 179-182.
- [16] Ellis, M.B. *Dematiaceous Hyphomycetes*. Kew, Surrey: Commonwealth Mycological Institute, 1971.
- [17] Ellis, M.B. *More Dematiaceous Hyphomycetes*. Kew, Surrey: Commonwealth Mycological Institute, 1976.
- [18] Gilman, J. "A Manual of Soil Fungi." Ames: Iowa State University Press, 1971.
- [19] Ramirez, C. "Manual and Atlas of Penicillia." Amsterdam: Elsevier Biomedical Press, 1982.
- [20] Raper, K.B. and Fennell, D.T. "The Genus *Aspergillus*." Baltimore, MD., Williams and Wilkins, Inc., 1965.
- [21] Buchanan, R.E. and Gibbons, N.E. *Bergey's Manual of Determinative Bacteriology*. Baltimore, Md: William and Wilkins, 1974.
- [22] Bower, C.A. and Wilcox, L.V. "Soluble Salts." In: *Methods of Soil Analysis*. Part 2: Black, C.A., Evan, D.D., White, J.L., Ensminger, L.E. and Clark, F.E. (Eds.), pp. 933-951., Madison: American Society of Agronomy, 1965.
- [23] Broadbent, F.E. "Organic Matter." In: *Methods of Soil Analysis*. Part 2: Black, C.S., Evan, D.D., White, J.L., Ensminger, L.E. and Clark, F.E. (Eds.), pp. 1397-1400. Madison: American Society of Agronomy, 1965.
- [24] Peech, M. "Hydrogen-ion Activity." In: *Methods of Soil Analysis*. Part 2: Black, C.S., Evan, D.D., White, J.L., Ensminger, L.E. and Clark, F.E. (Eds), pp. 914-925. Madison, American Society of Agronomy, 1965.
- [25] Piper, C.S. *Soils and Plant Analysis*. New York: Intersciences Publishers Inc., 1944.
- [26] Hashem, A.R. "Analysis of Water and Soils from Ashafa, Toroba, Wahat and Wehait." *J. King Saud University, Sc. 2*, No. 2 (1990), 87-94.
- [27] Hashem, A.R. "Soil Analysis and Mycoflora of the Industrial Yanbu City, Saudi Arabia." *Arab Gulf J. Scientific Research* 11, No. 1(1993), 91-103.

- [28] Hashem, A.R. "Heavy Metals Analysis of Water and Soils from Saudi Arabia." *J. King Saud University, Sci.* 5, No. 1 (1993), 39-46.
- [29] Bokhary, H. and Parvez, S. "Fungi from Petroleum-contaminated Saudi Arabian Soil." *Arid Soil Research and Rehabilitation* 7, (1993), 191-195.
- [30] Cerniglia, C.E. "Initial Reactions in the Oxidation of Anthracene by *Cunninghamella elegans*." *Journal of General Microbiology*, 128, (1982), 2055-2061.
- [31] Hettige, G. and Sheridan, J.E. "Interaction of Fungi Contaminating Diesel fuel." *Inter. Biodeter. Bull.* 25 (1984), 299-309.
- [32] Utkin, I.B., *et al.* "Degradation of Benzene, Toluene and Xylene by a *Pseudomonas* sp. 13 Culture." *Applied Biochemistry and Microbiology*, 28, No. 3 (1992), 275-280.
- [33] Harris, C.M. "Health Effects Research Initiatives at the Agency for Toxic Substances and Disease Registry-applicability to Contaminated Soils." In: *Petroleum Contaminated Soils*. Calabrese, E.J. and Kostecky, P.T. (Eds.), Vol. 2, pp. 49-53. Michigan: Lewis Publishing, Inc., 1990.

## أثر التلوث بالزيت الخام على الخواص الكيميائية والحيوية (الميكروبيولوجية) للتربة في المملكة العربية السعودية

عبدالوهاب رجب هاشم بن صادق

قسم النبات والأحياء الدقيقة، كلية العلوم، جامعة الملك سعود، ص. ب. ٢٤٥٥،

الرياض ١١٤٥١، المملكة العربية السعودية

(سُلم في ٢١ رجب ١٤١٤هـ؛ وقُبل للنشر في ١٤ شعبان ١٤١٥هـ)

ملخص البحث. تمَّ جمع عيّنات من التربة الملوّثة وغير الملوّثة بالبتروول من شرق المملكة العربية السعودية وحُللت لمعرفة قوام التربة والمحتوى العضوي الكلي للأملاح الذائبة وتفاعل التربة والمحتوى المعدني والفلورا الفطرية والبكتيرية.

أظهرت نتائج البحث أنّ التربة الملوّثة وغير الملوّثة رملية القوام معتدلة القلوية وتحتوي على الكمية نفسها من المادة العضوية والأملاح الكليّة الذائبة والمحتوى المعدني.

كما وجد أن عدد المستعمرات البكتيرية والفطرية في الجرام الواحد من التربة أعلى في التربة غير الملوّثة مقارنة بالتربة الملوّثة. وقد تمَّ عزل عشرين نوعاً من الفطريات تنتمي إلى تسعة أجناس من التربة غير الملوّثة بينما تمَّ عزل سبعة عشر نوعاً من التربة الملوّثة. والأجناس الفطرية السائدة هي: *اسبرجيليس* (سنة أنواع) يليه *بنيسليام* و*كلادوسبوريوم* (ثلاثة أنواع). أما الأجناس البكتيرية التي تمَّ عزلها فهي *ارثروباكترو باسيليس* و*ميكروكوكسي* و*بزيديموناس* و*ستابيلوكوكس*.